

early treatment with angiotensin converting enzyme inhibitors, β blockers, or both should be instituted. SRed and TRc can be used to classify boys with DMD as being at high or low risk. Selectively instituting early treatment and potentially slowing down the progress to DCM may be possible. SR may be used in future longitudinal studies assessing the beneficial effect of treatment of asymptomatic boys with DMD.

Tissue Doppler imaging echocardiography may be angle dependant. Chest deformity and the tachycardia made acquisition difficult. The high signal to noise ratio may be the greatest limitation of the technique with the current technology.

In this study we showed that SR and TV are significantly lower in asymptomatic boys with DMD, when conventional echocardiography failed to show any abnormality. Myocardial function in these patients appears to be impaired from a very young age. SR and TV can be reliably used to identify early myocardial dysfunction and predict an adverse outcome. These conclusions need further validation on independent datasets and larger population samples.

ACKNOWLEDGEMENTS

Dr N Giatrakos was supported by a grant from the Hellenic Heart Foundation (ELIKAR). Dr M Kinali was partly funded by the Dutch Parents project. The ongoing support of the Muscular Dystrophy Campaign to the Dubowitz Neuromuscular Centre is gratefully acknowledged. We thank Dr Eugenio Mercuri for his valuable contribution.

Authors' affiliations

N Giatrakos, D Dawson, P Nihoyannopoulos, Cardiology Department, NHLI, Imperial College, London, UK

M Kinali, F Muntoni, Department of Paediatrics, Dubowitz Neuromuscular Centre, Imperial College, London, UK

D Stephens, Department of Mathematics, Imperial College, London, UK

Correspondence to: Dr Nickolaos Giatrakos, Echocardiography Department, National Heart and Lung Institute, Imperial College, Hammersmith Hospital, Du Cane Road, London W12 0NN, UK; n.giatrakos@lycos.co.uk

Accepted 8 December 2005

REFERENCES

- 1 **Nigro G**, Comi U, Politano L, *et al.* The incidence and evolution of cardiomyopathy in Duchenne muscular dystrophy. *Int J Cardiol* 1990;**26**:271–7.
- 2 **Frankel KA**, Rosser RJ. The pathology of the heart in progressive muscular dystrophy: epimyocardial fibrosis. *Hum Pathol* 1976;**7**:375–86.
- 3 **Bushby K**, Muntoni F, Bourke P. 107th ENMC International Workshop: the management of cardiac involvement in muscular dystrophy and myotonic dystrophy. 7th–9th June 2002, Naarden, The Netherlands. *Neuromuscul Disord* 2003;**13**:166–72.
- 4 **Heimdal A**, Stoylen A, Torp H, *et al.* Real-time strain rate imaging of the left ventricle by ultrasound. *J Am Soc Echocardiogr* 1998;**11**:1013–9.
- 5 **Uematsu M**, Miyatake K, Tanaka N, *et al.* Myocardial velocity gradient as a new indicator of regional left ventricular contraction: detection by a two-dimensional tissue Doppler imaging technique. *J Am Coll Cardiol* 1995;**26**:217–23.

IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2005.077420

Atrial myocarditis: a possible cause of idiopathic enlargement of bilateral atria

In March 1997, a 72 year old woman with a cardiothoracic ratio of 90% on chest x ray was admitted to our hospital with a haemorrhagic gastric ulcer. However, at that time, she had no cardiac symptoms. Epstein's anomaly was excluded and no other congenital or acquired cardiac abnormality without atrial fibrillation and tricuspid regurgitation was detected. We then made the diagnosis of idiopathic dilatation of the bilateral atria. In October 2003, she came back to our hospital because of congestive heart failure with progressing mitral regurgitation. The echocardiogram revealed notably

enlarged atria (right atrium 115.0×73.7 mm in diameter; left atrium 102.0×53.5 mm in diameter) (panel A). However, no dilation of the ventricles was observed and systolic function was normal. Since bilateral heart failure was resistant to conventional treatment, she received surgical reduction of both atria and tricuspid and mitral valvular annuloplasty was performed because no organic valvular degeneration was confirmed in the operation. Microscopic examination revealed persistent myocarditis with severe scarring (panel B). The major histocompatibility complex (MHC) class

I antigens, which are required for the action of direct cytotoxic T cells, were found to be expressed on the myocytes and over expressed in the interstitial spaces (panel C). MHC class II antigens were also over-expressed in the interstitium. This case suggests that this congenital benign anomaly might be contributed to by atrial persistent myocarditis. This unique condition should be reinvestigated using up-to-date diagnostic tools.

M Habara, H Fujieda, Y Nakamura
ynakamura@mti.biglobe.ne.jp

